

Chapter 5

PREGNANCY

Table of Contents

	Page
Introduction	127
A. Carboxyhemoglobin Levels in Pregnancy	128
1. Fetal Blood	128
2. Carbon Monoxide Uptake During Smoking	129
3. Effect on the Fetus	130
4. Role of Nicotine	131
B. Association of Smoking with Birth Weight	131
1. Examination of Future Smokers Before Acquisition of the Smoking Habit	132
2. Identification of Chemical Factors Influencing the Fetus	133
C. Correlation of Human Studies and Animal Experiments	134
1. Variability of Carboxyhemoglobin Levels in Pregnant Women	134
2. Rabbit Experiments on Carbon Monoxide by Astrup et al. ...	135
3. Experiments by Other Investigators	136
D. Practical Aspects of Smoking in Pregnancy	136
Bibliography	138

1005105365

The reproductive system has been publicized as much as the cardiovascular and central nervous systems as target organs influenced by the carbon monoxide contained in cigarette smoke. Pregnant women are being warned that cigarette smoking causes low birth weight, prematurity and perinatal morbidity and mortality.

Although pregnant women who smoke have a higher incidence of low birth weight infants than nonsmokers, this association is not necessarily one of cause and effect. Since 1972, there have been additional epidemiologic studies which question a causal effect of smoking. Women who smoke are different from nonsmokers in their familial background, nutrition, socioeconomic factors and environmental exposures. The details of epidemiologic and environmental studies are reviewed in this chapter, as they relate to low birth weight. Mortality and morbidity statistics of the newborn have been excluded because they are influenced by factors which are remotely related to carbon monoxide contained in cigarette smoke.

1005105366

A. Carboxyhemoglobin Levels in Pregnancy

The blood levels of carboxyhemoglobin in pregnant women are in the same range of values as in nonpregnant women. For pregnant smokers, the range of values reported by various investigators is as follows:

<u>Investigators (year)</u>	<u>Number Pregnant Women</u>	<u>COHb % Saturation</u> mean \pm SD . . . (range)
Haddon <u>et al.</u> (1961) ¹	13	(0 - 8.4)
Heron (1962) ²	21	6.7 (1.6 - 14)
Young and Pugh (1963) ³	6	2.0 \pm 0.7 (1.0 - 3.0)
Younaszai <u>et al.</u> (1968) ⁴	16	8.3 (2.0 - 12.0)
Scoppetta (1968) ⁵	28	4.5 (0.5 - 15.0)
Baribaud <u>et al.</u> (1970) ⁶	27	4.4
Cole <u>et al.</u> (1972) ⁷	93	4.1 (0.5 - 14.0)

The above results show a wide range of values for carboxyhemoglobin in pregnant smokers. Although the mean values are higher than those seen for pregnant nonsmokers (omitted from above table), there is an overlapping in the values for both groups.

1005105367

1. Fetal Blood.

In most of the studies on maternal blood levels tabulated above, the corresponding fetal blood has been analyzed. The % saturation in the fetal blood is higher than the maternal blood because of the lower hemoglobin values in the former.

Significant!

2. Carbon Monoxide Uptake During Smoking.

Pavlou et al.⁸ obtained samples before and after smoking cigarettes.

The patients showed a steady rise from 1 - 4.2% in the control period to 5.3 - 11.2% at the end of the smoking period. The duration of the change was not followed. These transient changes in the maternal blood are probably not reflected in the fetal blood because of the slow diffusion of carbon monoxide through the placenta.⁹⁻¹¹ The rise in carboxyhemoglobin concentration in response to smoking a single cigarette was compared in pregnant women, pregnant women suffering from anemia, and nonpregnant women. Dow et al.¹² reported the following responses to smoking:

	% CO Hb		
	Before Smoking	After Smoking	Mean \pm S E Δ
pregnant (nonanemic)	1.68 \pm 0.27	5.55 \pm 0.34	3.87
pregnant anemic	1.68 \pm 0.2	6.7 \pm 0.2	5.0
nonpregnant (nonanemic)	2.24 \pm 0.2	4.3 \pm 0.19	2.11

The conclusion of Dow et al.¹² is that pregnant women, particularly those suffering from anemia, show a more-pronounced increase in carboxyhemoglobin levels than nonpregnant women. If the implication is that anemia causes an increase in uptake of carbon monoxide, that is fallacious as pointed out by Payne¹³ who calculated the absolute concentration from the percentage saturation.

The levels are as follows:

1005105368

	CO Hb in mg/100l		
	Before Smoking	After Smoking	Δ
pregnant (nonanemic)	203	672	469
pregnant anemic	150	596	446

In the above table, the uptake of carboxyhemoglobin as a result of smoking by anemic pregnant women is no greater than that for nonanemic subjects. When levels of carboxyhemoglobin are expressed in terms of absolute amounts, anemia does not increase the uptake of carbon monoxide in pregnant women.

3. Effect on the Fetus.

Asmussen and Kjeldsen¹⁴ have recently reported their observations on the ultrastructure of umbilical arteries from newborn children. They reported intimal changes seen in the arteries of children of smoking mothers which were not seen in those of nonsmoking mothers. They interpreted these differences as representing early signs of atherosclerosis, similar to those seen in rabbits exposed to carbon monoxide (see Chapter 3).

It should be pointed out that atherosclerosis is not a disease of the newborn and that the umbilical and placental arteries behave differently from the systemic arteries. Kawahara *et al.*¹⁵ have compared the response of human placenta and systemic blood vessels and noted the opposite effects of vasoconstrictors and vasodilators. Furthermore, the uteroplacental blood pool does not respond in a consistent manner to cigarette smoking. Cloeren *et al.*¹⁶ reported 5 pregnant women who initially showed a slight decrease followed by an increase whereas in the other five, there was only an increase in the

uteroplacental blood pool following cigarette smoking. This variability is consistent with the proposition that smokers are not a uniform population and that the placental vessels do not behave in a uniform fashion.

4. Role of Nicotine.

There is a change in the fetal breathing movements during maternal smoking. ¹⁷ Gennser et al. ¹⁷ explain the effect as a result of nicotine that has crossed the placental barrier. Although nicotine has been detected in samples of amniotic fluid, it is not uniformly seen in all smokers. Van Vunakis et al. ¹⁸ found nicotine in 22 out of 37 samples in smokers with the following wide distribution of concentration.

<u>No. of Smokers</u>	<u>Range of nicotine ng/ ml/amniotic fluid</u>
15	0 - 10
13	11 - 20
8	21 - 30
1	31 - 40

The wide range of nicotine concentration in the amniotic fluid reflects the variability of nicotine absorbed from cigarette smoke among smokers. It is unlikely that a dose response relationship can be derived from amniotic fluid levels and daily consumption of cigarettes.

B. Association of Smoking with Birth Weight

Until 1972, the epidemiologic studies indicated that smoking mothers had newborns of lower birth weight than nonsmoking mothers. The prevalent opinion expressed in the Surgeon General Reports (1967, 1969, 1971, 1972) was that smoking, particularly the elevated carboxyhemoglobin, was the cause of

1005105370

the low birth weight. The subject has been omitted from the 1973, 1974 and 1975 Reports so that recent events have not been publicized. They are as follows:

1. Examination of Future Smokers Before Acquisition of the Smoking Habit.

In 1972, Yerushalmy¹⁹ explored the question whether "this increase in low birth weight infants to smoking gravidas is due to the smoking or the smoker". Unlike previous epidemiologic studies, Yerushalmy investigated the reproductive performance of future smokers during the periods before they acquired the smoking habit. His conclusion was that women who subsequently became smokers had a high incidence of low birth weight infants during the period before they began to smoke. He questioned the role of smoking as an exogenous factor and proposed instead that the cause of the low birth weight lies in the smoker herself. The results of epidemiologic studies since 1972 have revealed the importance of maternal nutrition which, in turn, influences fetal nutrition.²⁰⁻²⁵ Prematurity and low birth weight are seen more frequently in the black group regardless of smoking habit.²⁶⁻²⁸ Socioeconomic factors, education, legitimacy, familial trends, ethnic origin and age of the mother influence the perinatal mortality.²⁹⁻³⁵ The role of smoking has not been evaluated because of the concurrent risk factors which differ between smokers and nonsmokers. The smoking habit of the father also influences the status of the newborn³⁶ suggesting that a genetic factor from both parents is transmitted to the offspring which, in turn, influences birth weight and perinatal morbidity and mortality. This subject is

1005105371

discussed further in Chapter 9. The original study of Yerushalmy in 1972,¹⁹ as well as the subsequent investigations cited above, have led to an exchange of correspondence among investigators,³⁷⁻⁵² The major criticism revealed in published letters is that epidemiologic studies cannot support causal relationships between smoking and low birth weight.

2. Identification of Chemical Factors Influencing the Fetus.

A reduction in birth weight from chronic carbon monoxide poisoning has not been reported. There are no published epidemiologic studies of women exposed occupationally to carbon monoxide. The literature on maternal poisoning indicates death of the fetus with carboxyhemoglobin levels as low as 10% saturation in the mother.⁵³⁻⁶¹ There is reason to suspect that this level was not the highest amount of carbon monoxide exposure causing fetal death.

In recent years, several chemical factors in the environment have been suspected of causing a reduction in birth weight. They include carbon monoxide and other air pollutants,⁶²⁻⁶⁴ plant constituents that are likely to be ingested,⁶⁵ high nitrate content in the water⁶⁶ and trace metals in the water, food and atmosphere.⁶⁷ Drugs such as salicylates,⁶⁸⁻⁷⁰ diphenylhydantions,⁷¹ caffeine,⁷² alcohol⁷²⁻⁷⁵ and hormones⁷⁶⁻⁸¹ have been reported to cause fetal malformations, prematurity and neonatal deaths. These chemical factors have not been considered in the epidemiologic investigation of risk factors (see above).

1005105372

C.: Correlation of Human Studies and Animal Experiments.

In 1972, Astrup *et al.*⁸² reported the results of human and animal studies and concluded that the "carbon monoxide in tobacco smoke might be responsible for the reduced birth weight of babies whose mothers smoked during pregnancy."

There are several features in this report which have been overlooked.

1. Variability of Carboxyhemoglobin Levels in Pregnant Women.

Pregnant women in the outpatient clinic of a hospital in Copenhagen were chosen at random and blood samples were taken for carboxyhemoglobin levels. The results indicate a considerable overlapping of values between smokers and nonsmokers:

COHb % Saturation	No. of Pregnant Women	
	Smokers	Nonsmokers
0-1%	336	678
1-2%	155	128
2-3%	139	51
3-4%	105	16
4-5%	50	7
5-6%	21	1
6-7%	11	2
7-8%	2	1
8-9%	4	0
9-10%	1	0
Total	824	884
Mean COHb	1.92%	0.87%

1005105373

Although the distribution of carboxyhemoglobin values was tabulated, only the mean values for birth weight were mentioned: 2990 g for smokers and 3225 g for nonsmokers. They further stated that, among smokers, there was a "negative correlation between birth weight and mean value of carboxyhemoglobin measurements of each individual."

Astrup et al.⁸² did not mention the correlation of birth weight with carboxyhemoglobin measurements of nonsmokers. In the discussion of their results, they were oblivious to the considerable literature on the role of risk factors, particularly of nutrition which was called to their attention by Neale.⁸³ Subsequent to 1972, the report of Yerushalmy¹⁹ and others appeared invalidating the premise of Astrup et al. that smoking was the cause of the low birth weight.

2. Rabbit Experiments on Carbon Monoxide by Astrup et al.

Astrup et al.⁸² exposed rabbits to carbon monoxide continuously for 30 days, from mating until the day before expected delivery. The rabbits that had concentrations of 16 to 18% had 35% stillbirths and offspring which died within the first hour, and lower birth weight than control rabbits. Since the levels of 16 to 18% were never attained in Astrup's patients, these experiments are irrelevant to the human studies.

The next exposure experiments by Astrup et al.⁸² consisted of rabbits exposed continuously to carboxyhemoglobin levels of 8 to 9 %. Again, this level was seen only in one out of 824 pregnant women. Astrup et al. did not

1005105374

report animals exposed to levels of 5% carboxyhemoglobin saturation, which would be relevant to their observations.

3. Experiments by Other Investigators.

Although Astrup et al.⁸² reported the results of their rabbit experiments in 1972, there has been no confirmation of their results. To the contrary, there are observations which question the relevance of carbon monoxide experiments to the smoking question. Kirschbaum et al.⁸⁴ injected nicotine and administered cigarette smoke to pregnant ewes. Gas exchange and cardiovascular function were unchanged in both the ewes and their fetuses. Chronic exposure to cigarette smoke in mice⁸⁵ and rats⁸⁶ that were pregnant did not influence birth weight nor induce malformations of fetuses. The experiments showing a reduction in birth weight of newborns of rats exposed to cigarette smoke are explained by a reduction in food intake by the pregnant rats.⁸⁷

D. Practical Aspects of Smoking in Pregnancy

That smoking during pregnancy causes an elevation in carboxyhemoglobin level and reduction in birth weight of the newborn is not supported by experimental and epidemiologic data and indeed is rejected by better-designed studies. Animals exposed to cigarette smoke do not duplicate the results of exposure to carbon monoxide. Furthermore, there is a considerable degree of overlapping of carboxyhemoglobin levels of smoking and nonsmoking women in pregnancy.

1005105375

Several risk factors in pregnancy have been identified as contributing to the low birth weight of infants of smokers, such as their familial background, nutritional status, socioeconomic factors and environmental exposure to chemicals.

1005105376

1. Haddon, W., Jr., Nesbitt, R.E.B., and Garcia, R., Smoking and pregnancy: carbon monoxide in blood during gestation and at term, Obstet. Gynecol., 18, 262-267, 1961.
2. Heron, H.J., The effects of smoking during pregnancy: A review with a preview, N.Z. Med. J., 61, 545-548, 1962.
3. Young, I.M., and Pugh, L.G.C.E., The carbon monoxide content of foetal and maternal blood, J. Obstet. Gynaecol., 70, 681-683, 1963.
4. Younoszai, M.K., Kacic, A., and Haworth, J.C., Cigarette smoking during pregnancy: The effect upon the hematocrit and acid-base balance of the newborn infant, Can. Med. Assoc. J., 99, 197-200, 1968.
5. Scoppetta, V., Sul contenuto di ossido di carbonio nel sangue circolante di gestanti fumatrici (On the carbon monoxide content of the circulating blood of pregnant smokers), Arch. Ostet. Ginecol., 73, 369-375, 1968.
6. Baribaud, L., Yacoub, M., Faure, J., Malinas, Y., and Cau, G., L'oxycarbonemie de L'enfant ne de mere fumeuse, (Blood carbon monoxide in children born of smoking mothers), Med. Legale Dommage Corporel, 3, 272-274, 1970.
7. Cole, P.V., Hawkins, L.H., and Roberts, D., Smoking during pregnancy and its effects on the fetus, J. Obstet. Gynaecol. Br. Commonw., 79, 782-787, 1972.
8. Pavlou, C., Chard, T., Landon, J., and Letchworth, A.T., Circulating levels of human placental lactogen in late pregnancy: the effect of glucose loading, smoking and exercise, Eur. J. Obstet. Gynecol. Reprod. Biol., 3, 45-49, 1973.
9. Longo, L.D., Carbon monoxide in the pregnant mother and fetus and its exchange across the placenta, Ann. N.Y. Acad. Sci., 174, 313-341, 1970.
10. Friberg, L., Nystrom, A., and Swanberg, H., Transplacental diffusion of carbon monoxide in human subjects, Acta. Physiol. Scand., 45, 363-368, 1959.
11. Gemzell, C.A., Robbe, H., and Strom, G., On the equilibration of carbon monoxide between human maternal and fetal circulation in vivo, Scand. J. Clin. Lab. Invest., 10, 372-378, 1958.

1005105377

12. Dow, T.G.B., Rooney, P.J., and Spence, M., Does anaemia increase the risks to the fetus caused by smoking in pregnancy?, Br. Med. J., 4, 253-254, 1975.
13. Payne, R.W., Anaemia, smoking, and pregnancy, Br. Med. J., 4, 521, 1975.
14. Asmussen, I., Kjeldsen, K., Intimal ultrastructure of human umbilical arteries. Observations on arteries from newborn children of smoking and non-smoking mothers, Circ. Res., 36, 579-589, 1975.
15. Kawahara, Y., and Yamada, S., Pharmacological studies on the vascular system of the human placenta, Nippon Yakurigaku Zasshi, 70, 207-214, 1974. (Japanese)
16. Cloeren, S.E., Lippert, T.H., and Fridrich, R., The influence of cigarette smoking on fetal heart rate and uteroplacental blood volume, Arch. Gynakol., 216, 15-22, 1974.
17. Gennser, G., Marsal, K., Brantmark, B., Maternal smoking and fetal breathing movements, Am. J. Obstet. Gynecol., 123, 861-867, 1975.
18. Van Vunakis, H., Langone, J.J., and Milunsky, A., Nicotine and cotinine in the amniotic fluid of smokers in the second trimester of pregnancy, Am. J. Obstet. Gynecol., 120, 64-66, 1974.
19. Yerushalmy, J., Infants with low birth weight born before their mothers started to smoke cigarettes, Am. J. Obstet. Gynecol., 112, 277-284, 1972.
20. Naeye, R.L., Blanc, W., and Paul, C., Effects of maternal nutrition on the human fetus, Pediatrics, 52, 494-503, 1973.
21. Miller, H.C., and Hassanein, K., Fetal malnutrition in white newborn infants: Maternal factors, Pediatrics, 52, 504-512, 1973.
22. Abdul-Karim, R.W. and Beydoun, S.N., Growth of the human fetus, Clin. Obstet. Gynecol., 17, 37-52, 1974.
23. Rush, D., Maternal nutrition during pregnancy in industrialized societies, Am. J. Dis. Child., 129, 430-434, 1975.

1005105378

24. Lechtig, A., Delgado, H., Lasky, R.E., Klein, R.E., Engle, P.L., Yarbrough, C., and Habicht, J.P., Maternal nutrition and fetal growth in developing societies, Am. J. Dis. Child., 129, 434-437, 1975.
25. Lechtig, A., Yarbrough, C., Delgado, H., Habicht, J.P., Martorell, R., and Klein, R.E., Influence of maternal nutrition on birth weight, Am. J. Clin. Nutr., 28, 1223-1233, 1975.
26. Miller, H.C., and Hassanein, K., Maternal factors in "fetally malnourished" black newborn infants, Am. J. Obstet. Gynecol., 118, 62-67, 1974.
27. Lubs, M.L.E., Racial differences in maternal smoking effects on the newborn infant, Am. J. Obstet. Gynecol., 115, 66-76, 1973.
28. Gendall, M., and Hellegers, A.E., The influence of the changes in maternal age, birth order, and color on the changing perinatal mortality, Baltimore, 1961-66, Health Serv. Rep., 88, 733-742, 1973.
29. Chabot, M.J., Garfinkel, J., and Pratt, M.W., Urbanization and differentials in white and nonwhite infant mortality, Pediatrics, 56, 777-781, 1975.
30. Grossman, S., Handlesman, Y., and Davies, A.M., Birth weight in Israel, 1968-70. I. Effects of birth order and maternal origin, J. Biosoc. Sci., 6, 43-58, 1974.
31. Handlesman, Y., and Davies, A.M., Birth weight in Israel, 1968-70. II. The effect of paternal origin, J. Biosoc. Sci., 7, 153-164, 1975.
32. Bjerre, I., and Varendh, G., A study of some biological and socioeconomic factors in low birth weight, Acta Paediatr. Scand., 64, 605-612, 1975.
33. Shin, E.H., Black-white differentials in infant mortality in the south, 1940-1970, Demography, 12, 1-19, 1975.
34. Dott, A.B., and Fort, A.T., The effect of maternal demographic factors on infant mortality rates. Summary of the findings of the Louisiana infant mortality study. Part I., Am. J. Obstet. Gynecol., 123, 847-853, 1975.

1005105379

35. Johnstone, F., and Inglis, L., Familial trends in low birth weight, Br. Med. J., 3, 659-661, 1974.
36. Mau, G., and Netter, P., The effects of paternal cigarette smoking on perinatal mortality and the incidence of malformations, Dtsch. Med. Wochenschr., 99, 1113-1118, 1974.
37. Yerushalmy, J., Cigarette smoking, infant birth weight, and perinatal mortality rates, Am. J. Obstet. Gynecol., 118, 884-886, 1974.
38. Williams, H.S., and Meyer, M.B., Reply to Dr. Yerushalmy, Am. J. Obstet. Gynecol., 118, 886-888, 1974.
39. Yerushalmy, J., Smoking in pregnancy, Dev. Med. Child Neurol., 15, 691-693, 1973.
40. Ross, E.M., Butler, N.R., and Goldstein, H., Smoking hazards to the fetus, Br. Med. J., 4, 51, 1973.
41. Burch, P.R.J., Smoking hazard to the fetus, Br. Med. J., 1, 40, 1974.
42. Ounsted, M., Familial trends in low birth weight, Br. Med. J., 4, 163, 1974.
43. James, W.H., Smoking in pregnancy and child development, Br. Med. J., 2, 610-611, 1974.
44. Yerushalmy, J., Congenital heart disease and maternal smoking habits, Nature, 242, 262, 1973.
45. Fedrick, J., Alberman, E.D., and Goldstein, H., Reply to professor Yerushalmy's letter on congenital heart disease, Nature, 242, 263, 1973.
46. Burch, P.R.J., Smoking, pregnancy, and publicity, Nature, 245, 277, 1973.
47. Goldstein, H., Smoking pregnancy and publicity, Nature, 245, 467-468, 1973.
48. Bruch, P.R.J., Smoking and pregnancy, Nature, 246, 177, 1973.
- 48a. Hickey, R.J., Boyce, D.E., and Harner, E.B., Smoking and pregnancy, Nature, 246, 177-178, 1973.

1005105380

49. James, W.H., Smoking in pregnancy, Nature, 246, 235, 1973.
50. Goldstein, H., Smoking, pregnancy and publicity, Nature, 246, 540, 1973.
51. Goldstein, H., Smoking and pregnancy, Nature, 245, 277, 1973.
52. Hardy, J.B., and Mellits, E.D., Effect on child of maternal smoking during pregnancy, Lancet, 1, 719-720, 1973.
53. Goldstein, D.P., Carbon monoxide poisoning in pregnancy, Am. J. Obstet. Gynecol., 92, 526-528, 1965.
54. Kamraj-Mazurkiewicz, K., Effect of carbon monoxide poisoning in the course of pregnancy on the fetal central nervous system according to our case, Ginek. Polska, 38, 291-294, 1967. (Polish)
55. Bankl, H., and Jellinger, K., Central nervous system injuries following fetal carbon monoxide poisoning, Beitr. Pathol. Anat., 135, 350-376, 1967. (German)
56. Szilagyi, L., Chronic neurologic damage from fetal CO poisoning, Orv. Hetil., 108, 1559-1561, 1967. (Hungarian)
57. Kells, D.U. Carbon monoxide poisoning during pregnancy, Med. Leg. Bull., 177, 1-11, 1968.
58. Beaudoin, A., Gachon, J., Butin, L.P., and Bost, M., Fetal consequences of carbon monoxide poisoning of the mother, Pediatric, 24, 539-553, 1969. (French)
59. Stokowski, C., and Keskak, J., Carbon monoxide poisoning in pregnancy, Ginek. Pol., 40, 801-804, 1969. (Polish)
60. Matsuyama, H., Influence of maternal fuel-gas poisoning upon fetal brain, Adv. Neurol. Sci. Tokyo, 34-38, 1969. (Japanese)
61. Larcan, A., Landes, P., and Vert, P., Carbon monoxide during the 2nd month of pregnancy without neonatal abnormality, Bull. Fed. Soc. Gynecol. Obstet. Lang. Fr., 22, 338-339, 1970. (French)

1005105381

62. Malinas, Y., Health care in pregnancy, Concours Med., 96, 45-48, 1974. (French)
63. Williams, L.A., The effect of air pollution on human birth weight, Am. J. Phys. Anthropol., 42, 340, 1975.
64. Hunt, V.R., and Cross, W.L., Infant mortality and the environment of a lesser metropolitan county: A study based on births in one calendar year, Environ. Res., 9, 135-151, 1975.
65. James, L.F., Plants - natural sources of teratogenicity, Clin. Toxicol., 6, 297-298, 1973.
66. Gelperin, A., Moses, V.K., and Bridger, C., Relationship of high nitrate community water supply to infant and fetal mortality, Ill. Med. J., 155-157 & 186, February, 1975.
67. Baglan, R.J., Brill, A.B., Schulert, A., Wilson, D., Larsen, K., Dyer, N., Mansour, M., Schaffner, W., Hoffman, L., and Davies, J., Utility of placental tissue as an indicator of trace element exposure to adult and fetus, Environ. Res., 8, 64-70, 1974.
68. Collins, E., and Turner, G., Maternal effects of regular salicylate ingestion in pregnancy, Lancet, 2, 335-337, 1975.
69. Turner, G., and Collins, E., Fetal effects of regular salicylate ingestion in pregnancy, Lancet, 2, 338-339, 1975.
70. Collins, E., and Turner, G., Salicylates and pregnancy, Lancet, 2, 1494, 1973.
71. Monson, R.R., Rosenberg, L., Hartz, S.C., Shapiro, S., Heinonen, O. P., and Slone, D., Diphenylhydantoin and selected congenital malformations, N. Engl. J. Med., 289, 1049-1052, 1973.
72. Mau, G., Subsistence and luxury foods consumption during pregnancy and its effect on perinatal mortality, prematurity, and other perinatal factors, Monatsschr. Kinderheilkd., 122, 539-540, 1974. (German)
73. Anonymous, Defects studied in infants of alcoholic mothers, J. Am. Med. Assoc., 226, 520-521, 1973.

1005105382

74. Jones, K.L., and Smith, D.W., Recognition of the fetal alcohol syndrome in early infancy, Lancet, 2, 999-1001, 1973.
75. Tenbrinck, M.S., and Buchin, S.Y., Fetal alcohol syndrome, Report of a case, J. Am. Med. Assoc., 232, 1144-1147, 1975.
76. Mercke, C., Cavallin-Stahl, E., and Lundh, B., Carbon monoxide production and reticulocyte count in normal women, Acta Med. Scand., 198, 155-160, 1975.
77. Beischer, N.A., Low oestriol excretion incidence, significance and treatment in an obstetric population, Med. J. Aust., 2, 379-381, 1975.
78. Canick, J.A., and Ryan, K.J., The inhibition of 16 α -hydroxytestosterone aromatization by carbon monoxide, Biochem. Biophys. Res. Commun., 63, 496-501, 1975.
79. Johnson, J.W.C., Austin, K.L., Jones, G.S., Davis, G.H., King, T.M., Efficacy of 17 α -hydroxprogesterone caproate in the prevention of premature labor, N. Engl. J. Med., 293, 675-680, 1975.
80. Drew, J.H., Goldenberg, H.J., and Beischer, N.A., Children born from pregnancies complicated by subnormal oestriol excretion - first year, Med. J. Aust., 2, 381-384, 1975.
81. Abell, D.A., and Beischer, N.A., The influence of abnormal glucose tolerance (hyperglycaemia and hypoglycaemia) on pregnancy outcome when oestriol excretion is subnormal, Br. J. Obstet. Gynaecol., 82, 936-942, 1975.
82. Astrup, P., Olsen, H.M., Trolle, D., and Kjeldsen, K., Effect of moderate carbon-monoxide exposure on fetal development, Lancet, 2, 1220-1222, 1972.
83. Neale, R.J., Effect of moderate carbon-monoxide exposure on fetal development, Lancet, 2, 1419, 1972.
84. Kirschbaum, T.H., Dilts, P.V., Jr., and Brinkman, C.R., III, Some acute effects of smoking in sheep and their fetuses, Obstet. Gynecol., 35, 527-536, 1970.

1005105383

85. Wagner, B., Lazar, Ph., and Chouroulinkov, I., The effects of cigarette smoke inhalation upon mice during pregnancy, Rev. Eur. Etud. Clin. Biol., 17, 943-948, 1972.
86. Reckzeh, G., Dontenwill, W., and Leuschner, F., Testing of cigarette smoke inhalation for teratogenicity in rats, Toxicology, 4, 289-295, 1975.
87. Younoszai, M.K. Peloso, J., and Haworth, J.C. Fetal growth retardation in rats exposed to cigarette smoke during pregnancy, Am. J. Obstet. Gynecol., 104, 1207-1213, 1969.

1005105384